



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES**  
(Untabulated).

**ALGERIA—Algiers.**—Month of February, 1912. Population, 172,397. Total number of deaths from all causes 320, including diphtheria 4, tuberculosis 48, typhoid fever 5.

**BRITISH BURMA—Rangoon.**—Month of December, 1911. Population, 293,316. Total number of deaths from all causes, 811, including measles 1, plague 11, smallpox 8, tuberculosis 56, typhoid fever 5.

**FRANCE—Calais.**—Month of February, 1912. Population, 80,000. Total number of deaths from all causes 118, including measles 1, scarlet fever 2, tuberculosis 22.

**GREAT BRITAIN.**—Week ended February 24, 1912.

*England and Wales.*—The deaths registered in 77 great towns correspond to an annual rate of 16.8 per 1,000 of the population, which is estimated at 17,559,219.

*Ireland.*—The deaths registered in 21 principal town districts correspond to an annual rate of 24.6 per 1,000 of the population, which is estimated at 1,157,014. The lowest rate was recorded at Armagh, viz, 6.9, and the highest at Galway, viz, 43.3, per 1,000.

*Scotland.*—The deaths registered in 18 principal towns correspond to an annual rate of 18.5 per 1,000 of the population, which is estimated at 2,182,400. The lowest rate was recorded at Falkirk, viz, 6.1, and the highest at Greenock, viz, 23.4, per 1,000. The total number of deaths from all causes was 775, including diphtheria 22, measles 62, scarlet fever 3, typhoid fever 4.

**ITALY—Genoa.**—Two weeks ended February 29, 1912. Population 272,077. Total number of deaths from all causes, 192, including diphtheria 7, measles 1, scarlet fever 1, tuberculosis 56, typhoid fever 1.

*Milan.*—Month of February, 1912. Population 602,236. Total number of deaths from all causes 138, including diphtheria 1, measles 6, tuberculosis 120, typhoid fever 8.

**SPAIN—Almeria.**—Month of February, 1912. Population 50,910. Total number of deaths from all causes 123, including tuberculosis 14, typhoid fever 3.

*Madrid.*—Month of February, 1912. Population 584,117. Total number of deaths from all causes 1,213, including diphtheria 12, measles 9, tuberculosis 123, typhoid fever 22.

*Seville.*—Month of February, 1912. Population 158,235. Total number of deaths from all causes 386, including diphtheria 5, smallpox 3, tuberculosis 69, typhoid fever 5.

TAHITI.—Four weeks ended February 16, 1912. Population 4,000. Total number of deaths from all causes 5. No deaths from contagious diseases.

By authority of the Secretary of the Treasury:

RUPERT BLUE,  
*Surgeon General,*  
*United States Public Health and Marine-Hospital Service.*

---

---

ADDITIONAL COPIES of this publication  
may be procured from the SUPERINTEND-  
ENT OF DOCUMENTS, Government Printing  
Office, Washington, D. C., at 5 cents per copy.

Subscription price, per volume ..... \$2

---

